



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GEOMETRY.

384. Proposed by S. LEFSEHETZ, Clark University.

Let ABC be a triangle, O a circle tangent to its three sides, T a variable tangent of O , which cuts the sides BC , CA , AB in a , b , c . Oa' , Ob' , Oc' the perpendiculars in O to Oa , Ob , Oc , cutting, respectively, T in points a' , b' , c' . Prove that Aa' , Bb' , Cc' meet in a point t , and find the locus of t when T varies. Purely geometrical proofs wanted.

385. Proposed by V. M. SPUNAR, M. and E. E., Chicago, Ill.

Given a triangle ABC , find the radius of a circle touching two of its sides and a line parallel to the third, at a distance $d=u+2r$.

386. Proposed by DANIEL KRETH, Oxford, Iowa.

Construct the triangle, having given, the vertical angle, the sum of the three sides, and the perpendicular.

CALCULUS.

308. Proposed by C. N. SCHMALL, New York City.

Prove, by calculus, that of all isoperimetric triangles, the equilateral has the greatest area.

309. Proposed by S. G. BARTON, Ph. D., Clarkson School of Technology.

In practical problems involving maxima and minima, it is really the greatest and least values of the function which are desired. Show why we can assume that the maximum is the greatest value and the minimum the least value under the conditions.

310. Proposed by C. N. SCHMALL, New York City.

Evaluate $\int_0^\pi \frac{dx}{1-2a\cos x + a^2}$. Edwards' *Integral Calculus for Beginners*, page 131, ex. 9, (iii). The answer given is $\frac{\pi}{1-a^2}$. Is this a complete answer to the question?

MECHANICS.

260. Proposed by W. J. GREENSTREET, M. A., Stroud, England.

To the ends of a fine inextensible string, length $2l$, are attached to equal, smooth, spherical, equally elastic (e) particles. At first the middle point of the string touches a rigid, fixed, circular rim, radius a , and the particles are $2l$ apart. They are now projected with equal velocities perpendicular to the string and curl around the rim. If l is greater than πa , find the condition that the particles will move after collision along tangents to the rim, the whole motion being on a smooth horizontal plane.

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

183. Proposed by M. T. GOODRICH, Dixfield, Maine.

Show what relation must exist between the quantities A , B , and C , in the harmonic ratio $\frac{AB}{(A+B+C)(-C)} = -1$, so that they will be real positive integers.